

Rivermead Post Concussion Symptoms Questionnaire

Modified (Rpq-3 And Rpq-13)⁴² Printed With Permission: Modified Scoring System From Eyres 2005 ²⁸

Name:

Date:

After a head injury or accident some people experience symptoms that can cause worry or nuisance. We would like to know if you now suffer any of the symptoms given below. Because many of these symptoms occur normally, we would like you to compare yourself now with before the accident. For each symptom listed below please circle the number that most closely represents your answer.

0 = not experienced at all
 1 = no more of a problem
 2 = a mild problem
 3 = a moderate problem
 4 = a severe problem

Compared with **before** the accident, do you **now** (i.e., over the last 24 hours) suffer from:

	not experienced	no more of a problem	mild problem	moderate problem	severe problem
Headaches	0	1	2	3	4
Feelings of dizziness	0	1	2	3	4
Nausea and/or vomiting	0	1	2	3	4
Noise sensitivity (easily upset by loud noise)	0	1	2	3	4
Sleep disturbance	0	1	2	3	4
Fatigue, tiring more easily	0	1	2	3	4
Being irritable, easily angered	0	1	2	3	4
Feeling depressed or tearful	0	1	2	3	4
Feeling frustrated or impatient	0	1	2	3	4
Forgetfulness, poor memory	0	1	2	3	4
Poor concentration	0	1	2	3	4
Taking longer to think	0	1	2	3	4
Blurred vision	0	1	2	3	4
Light sensitivity (easily upset by bright light)	0	1	2	3	4
Double vision	0	1	2	3	4
Restlessness	0	1	2	3	4

Are you experiencing any other difficulties? Please specify, and rate as above.

1.	0	1	2	3	4
2.	0	1	2	3	4

Administration only:

RPQ-3 (total for first three items)	
RPQ-13 (total for next 13 items)	

Rivermead Post Concussion Symptoms Questionnaire (cont.)

Modified (Rpq-3 And Rpq-13)⁴² Printed With Permission: Modified Scoring System From Eyres 2005 ²⁸

Administration only

Individual item scores reflect the presence and severity of post concussive symptoms. Post concussive symptoms, as measured by the RPQ, may arise for different reasons subsequent to (although not necessarily directly because of) a traumatic brain injury. The symptoms overlap with broader conditions, such as pain, fatigue and mental health conditions such as depression⁷².

The questionnaire can be repeated to monitor a patient's progress over time. There may be changes in the severity of symptoms, or the range of symptoms. Typical recovery is reflected in a reduction of symptoms and their severity within three months.

Scoring

The scoring system has been modified from Eyres, 2005²⁴.

The items are scored in two groups. The first group (RPQ-3) consists of the first three items (headaches, feelings of dizziness and nausea) and the second group (RPQ-13) comprises the next 13 items. The total score for RPQ-3 items is potentially 0–12 and is associated with early symptom clusters of post concussive symptoms. If there is a higher score on the RPQ-3, earlier reassessment and closer monitoring is recommended.

The RPQ-13 score is potentially 0–52, where higher scores reflect greater severity of post concussive symptoms. The RPQ-13 items are associated with a later cluster of symptoms, although the RPQ-3 symptoms of headaches, dizziness and nausea may also be present. The later cluster of symptoms is associated with having a greater impact on participation, psychosocial functioning and lifestyle. Symptoms are likely to resolve within three months. A gradual resumption of usual activities is recommended during this period, appropriate to symptoms. If the symptoms do not resolve within three months, consideration of referral for specialist assessment or treatment services is recommended.

References:

Eyres, S., Carey, A., Gilworth, G., Neumann, V., Tennant, A. (2005). Construct validity and reliability of the Rivermead Post Concussion Symptoms Questionnaire. *Clinical Rehabilitation*, 19, 878-887.

King, N. S., Crawford, S., Wenden, F.J., Moss, N.E.G. Wade, D.T. (1995). The Rivermead Post Concussion Symptoms Questionnaire: a measure of symptoms commonly experienced after head injury and its reliability *Journal of Neurology*, 242, 587-592.

Potter, S., Leigh, E., Wade, D., Fleminger, S. (2006). The Rivermead Post Concussion Symptoms Questionnaire *Journal of Neurology*, October 1-12.